

Specimen ID:
Control ID:

Acct #: 17452095
Walk-In Lab, LLC
VART verified
169 W Augusta Lane
SLIDELL, LA 70458

Phone: (800) 539-6119 Rte: 00

Patient Details
DOB:
Age(y/m/d):
Gender: SSN:
Patient ID:

Specimen Details
Date collected:
Date entered:
Date reported:

Physician Details
Ordering: F AN
Referring:
ID: 1558345843
NPI: 1558345843

General Comments & Additional Information

Alternate Control Number:
Total Volume: Not Provided

Alternate Patient ID: Not Provided
Fasting: No

Ordered Items

CBC With Differential/Platelet; Testosterone,Free and Total; Prostate-Specific Ag, Serum; Venipuncture

TESTS	RESULT	FLAG	UNITS	REFERENCE	INTERVAL	LAB
CBC With Differential/Platelet						
WBC	13.2	High	x10E3/uL	3.4 - 10.8		01
RBC	5.21		x10E6/uL	4.14 - 5.80		01
Hemoglobin	17.2		g/dL	12.6 - 17.7		01
Hematocrit	49.9		%	37.5 - 51.0		01
MCV	96		fL	79 - 97		01
MCH	33.0		pg	26.6 - 33.0		01
MCHC	34.5		g/dL	31.5 - 35.7		01
RDW	13.6		%	12.3 - 15.4		01
Platelets	275		x10E3/uL	150 - 379		01
Neutrophils	65		%			01
Lymphs	23		%			01
Monocytes	11		%			01
Eos	1		%			01
Basos	0		%			01
Neutrophils (Absolute)	8.6	High	x10E3/uL	1.4 - 7.0		01
Lymphs (Absolute)	3.1		x10E3/uL	0.7 - 3.1		01
Monocytes(Absolute)	1.4	High	x10E3/uL	0.1 - 0.9		01
Eos (Absolute)	0.1		x10E3/uL	0.0 - 0.4		01
Baso (Absolute)	0.0		x10E3/uL	0.0 - 0.2		01
Immature Granulocytes	0		%			01
Immature Grans (Abs)	0.0		x10E3/uL	0.0 - 0.1		01
Testosterone,Free and Total						
Testosterone, Serum	6579	High	ng/dL	348 - 1197		01
Results confirmed on dilution.						
Comment :						
Adult male reference interval is based on a population of lean males up to 40 years old.						
Free Testosterone(Direct)	>50.0	High	pg/mL	6.8 - 21.5		02

Patient:
DOB:

Patient ID:

Control ID:

Specimen ID:
Date collected:

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
-------	--------	------	-------	--------------------	-----

Prostate-Specific Ag, Serum

Prostate Specific Ag, Serum	1.1		ng/mL	0.0 - 4.0	01
-----------------------------	-----	--	-------	-----------	----

Roche ECLIA methodology.

According to the American Urological Association, Serum PSA should decrease and remain at undetectable levels after radical prostatectomy. The AUA defines biochemical recurrence as an initial PSA value 0.2 ng/mL or greater followed by a subsequent confirmatory PSA value 0.2 ng/mL or greater.

Values obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.

01	SO	LabCorp San Diego 13112 Evening Creek Dr So Ste 200, San Diego, CA 92128-4108	Dir: Jenny Galloway, MD
02	BN	LabCorp Burlington 1447 York Court, Burlington, NC 27215-3361	Dir: William F Hancock, MD

For inquiries, the physician may contact **Branch: 504-828-2666 Lab: 858-668-3700**