

Patient Information	Specimen Information	Client Information
DOB: AGE: Gender: Phone: Patient ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	

Test Name	In Range	Out Of Range	Reference Range	Lab
TSI (THYROID STIMULATING IMMUNOGLOBULIN) TSI	<89		<140 % baseline	

Thyroid stimulating immunoglobulins (TSI) can engage the TSH receptors resulting in hyperthyroidism in Graves' disease patients. TSI levels can be useful in monitoring the clinical outcome of Graves' disease as well as assessing the potential for hyperthyroidism from maternal-fetal transfer. TSI results greater than or equal to (>=) 140% of the Reference Control are considered positive.

NOTE: A serum TSH level greater than 350 micro-International Units/mL can interfere with the TSI bioassay and potentially give false positive results.

Patients who are pregnant and are suspected of having hyperthyroidism should have both a TSI and human Chorionic Gonadotropin (hCG) tests measured. A serum hCG level greater than 40,625 mIU/mL can interfere with the TSI bioassay and may give false negative results. In these patients it is recommended that a second TSI is obtained when the hCG concentration falls below 40,625 mIU/mL (usually after approximately 20-weeks gestation).

PERFORMING SITE: