

Patient Information	Specimen Information	Client Information
DOB: AGE: Gender: Phone: Patient ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	

COMMENTS: FASTING:NO

Test Name	In Range	Out Of Range	Reference Range	Lab
URIC ACID Therapeutic target for gout patients: <6.0 mg/dL		7.1 H	2.5-7.0 mg/dL	
SED RATE BY MODIFIED WESTERGREN RHEUMATOID FACTOR C-REACTIVE PROTEIN	<14 5.9	39 H	< OR = 20 mm/h <14 IU/mL <8.0 mg/L	

Patient Information	Specimen Information	Client Information
DOB: AGE: Gender: Patient ID:	Specimen: Collected: Received: Reported:	

Immunology

Test Name	Result	Reference Range	Lab
-----------	--------	-----------------	-----

ANA SCREEN, IFA, W/REFL TITER AND PATTERN

ANA SCREEN, IFA	NEGATIVE	NEGATIVE	
-----------------	----------	----------	--

ANA IFA is a first line screen for detecting the presence of up to approximately 150 autoantibodies in various autoimmune diseases. A negative ANA IFA result suggests ANA-associated autoimmune diseases are not present at this time.

Visit Physician FAQs for interpretation of all antibodies in the Cascade, prevalence, and association with diseases at <http://education.QuestDiagnostics.com/faq/FAQ177>

Physician Comments:

PERFORMING SITE: