

Patient Information	Specimen Information	Client Information
DOB: AGE: Gender: Phone: Patient ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	

COMMENTS: FASTING:NO

Test Name In Range Out Of Range Reference Range Lab

ALPHA FETOPROTEIN,

TUMOR MARKER 2.

2.5 ng/mL

Reference Range: <6.1

The use of AFP as a tumor marker in pregnant females is not recommended.

This test was performed using the Beckman Coulter chemiluminescent method. Values obtained from different assay methods cannot be used interchangeably. AFP levels, regardless of value, should not be interpreted as absolute evidence of the presence or absence of disease.

CEA <0.5 ng/mL

Non-Smoker: <2.5 Smoker: <5.0

This test was performed using the Siemens chemiluminescent method. Values obtained from different assay methods cannot be used interchangeably. CEA levels, regardless of value, should not be interpreted as absolute evidence of the presence or absence of disease.

CA 19-9 11 <34 U/mL

This test was performed using the Siemens chemiluminescent method. Values obtained from different assay methods cannot be used interchangeably. CA 19-9 levels, regardless of value, should not be interpreted as absolute evidence of the presence or absence of disease.

CA 15-3 5 <32 U/mL

This test was performed using the Siemens (Bayer) chemiluminescent method. Values obtained from different assay methods cannot be used interchangeably. CA 15-3 levels, regardless of value, should not be interpreted as absolute evidence of the presence or absence of disease.

CA 125 10 <35 U/mL

This test was performed using the Beckman Coulter Chemiluminescent method. Values obtained from different assay methods cannot be used interchangeably. CA 125 levels, regardless of value, should not be interpreted as absolute evidence of the presence or absence of disease.



Patient Information	Specimen Information	Client Information
DOB: AGE:	Specimen: Collected: Received: Reported:	

PERFORMING SITE: