

Patient Information	Specimen Information	Client Information
<b>DOB:</b> <b>AGE:</b> Gender: Phone: Patient ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	

**COMMENTS:**      FASTING: YES

Test Name	In Range	Out Of Range	Reference Range	Lab
PROTHROMBIN W/INR + PARTIAL THROMBOPLASTIN TIMES				
PARTIAL THROMBOPLASTIN TIME, ACTIVATED	27		22-34 sec	
This test has not been validated for monitoring unfractionated heparin therapy. For testing that is validated for this type of therapy, please refer to the Heparin Anti-Xa assay (test code 30292).  For additional information, please refer to <a href="http://education.QuestDiagnostics.com/faq/FAQ159">http://education.QuestDiagnostics.com/faq/FAQ159</a> (This link is being provided for informational/educational purposes only.)				
PROTHROMBIN TIME-INR				
INR	1.0			
Reference Range		0.9-1.1		
Moderate-intensity Warfarin Therapy		2.0-3.0		
Higher-intensity Warfarin Therapy		3.0-4.0		
PT	10.5		9.0-11.5 sec	
<b>HS CRP</b>		<b>3.6 H</b>	mg/L	
Higher relative cardiovascular risk according to AHA/CDC guidelines. Consider retesting in 1 to 2 weeks to exclude a benign transient elevation in the baseline CRP value secondary to infection or inflammation.  For ages >17 Years: hs-CRP mg/L    Risk According to AHA/CDC Guidelines <1.0                Lower relative cardiovascular risk. 1.0-3.0            Average relative cardiovascular risk. 3.1-10.0          Higher relative cardiovascular risk. Consider retesting in 1 to 2 weeks to exclude a benign transient elevation in the baseline CRP value secondary to infection or inflammation. >10.0             Persistent elevation, upon retesting, may be associated with infection and inflammation.				
COMPREHENSIVE METABOLIC PANEL				
GLUCOSE	82		65-99 mg/dL	
Fasting reference interval				
UREA NITROGEN (BUN)	11		7-25 mg/dL	
CREATININE	0.81		0.50-1.05 mg/dL	
For patients >49 years of age, the reference limit for Creatinine is approximately 13% higher for people identified as African-American.				

Patient Information	Specimen Information	Client Information
<b>DOB:</b> <b>AGE:</b> Gender: Patient ID:	Specimen: Collected: Received: Reported:	

Test Name	In Range	Out Of Range	Reference Range	Lab
eGFR NON-AFR. AMERICAN	85		> OR = 60 mL/min/1.73m2	
eGFR AFRICAN AMERICAN	98		> OR = 60 mL/min/1.73m2	
BUN/CREATININE RATIO	NOT APPLICABLE		6-22 (calc)	
SODIUM	139		135-146 mmol/L	
POTASSIUM	4.2		3.5-5.3 mmol/L	
CHLORIDE	104		98-110 mmol/L	
CARBON DIOXIDE	24		20-32 mmol/L	
CALCIUM	9.6		8.6-10.4 mg/dL	
PROTEIN, TOTAL	7.3		6.1-8.1 g/dL	
ALBUMIN	4.7		3.6-5.1 g/dL	
GLOBULIN	2.6		1.9-3.7 g/dL (calc)	
ALBUMIN/GLOBULIN RATIO	1.8		1.0-2.5 (calc)	
BILIRUBIN, TOTAL	0.5		0.2-1.2 mg/dL	
ALKALINE PHOSPHATASE	50		33-130 U/L	
AST	18		10-35 U/L	
ALT	19		6-29 U/L	
SED RATE BY MODIFIED WESTERGREN	6		< OR = 20 mm/h	
CBC (INCLUDES DIFF/PLT)				
WHITE BLOOD CELL COUNT	8.3		3.8-10.8 Thousand/uL	
RED BLOOD CELL COUNT	5.03		3.80-5.10 Million/uL	
HEMOGLOBIN	14.7		11.7-15.5 g/dL	
HEMATOCRIT	43.9		35.0-45.0 %	
MCV	87.3		80.0-100.0 fL	
MCH	29.2		27.0-33.0 pg	
MCHC	33.5		32.0-36.0 g/dL	
RDW	11.8		11.0-15.0 %	
PLATELET COUNT	227		140-400 Thousand/uL	
MPV	11.4		7.5-12.5 fL	
ABSOLUTE NEUTROPHILS	5445		1500-7800 cells/uL	
ABSOLUTE LYMPHOCYTES	2100		850-3900 cells/uL	
ABSOLUTE MONOCYTES	315		200-950 cells/uL	
ABSOLUTE EOSINOPHILS	349		15-500 cells/uL	
ABSOLUTE BASOPHILS	91		0-200 cells/uL	
NEUTROPHILS	65.6		%	
LYMPHOCYTES	25.3		%	
MONOCYTES	3.8		%	
EOSINOPHILS	4.2		%	
BASOPHILS	1.1		%	
URINALYSIS, COMPLETE				
COLOR	YELLOW		YELLOW	
APPEARANCE	CLEAR		CLEAR	
SPECIFIC GRAVITY	1.007		1.001-1.035	
PH	6.5		5.0-8.0	
GLUCOSE	NEGATIVE		NEGATIVE	
BILIRUBIN	NEGATIVE		NEGATIVE	
KETONES	NEGATIVE		NEGATIVE	
OCCULT BLOOD	NEGATIVE		NEGATIVE	
PROTEIN	NEGATIVE		NEGATIVE	
NITRITE	NEGATIVE		NEGATIVE	
<b>LEUKOCYTE ESTERASE</b>		<b>3+</b>	NEGATIVE	
<b>WBC</b>		<b>6-10</b>	< OR = 5 /HPF	
RBC	0-2		< OR = 2 /HPF	
SQUAMOUS EPITHELIAL CELLS	0-5		< OR = 5 /HPF	
<b>BACTERIA</b>		<b>FEW</b>	NONE SEEN /HPF	

Patient Information	Specimen Information	Client Information
<b>DOB:</b> <b>AGE:</b> Gender: Patient ID:	Specimen: Collected: Received: Reported:	

Test Name	In Range	Out Of Range	Reference Range	Lab
HYALINE CAST	NONE SEEN		NONE SEEN /LPF	
RHEUMATOID FACTOR	<14		<14 IU/mL	

Walk-In Lab

Patient Information	Specimen Information	Client Information
<b>DOB:</b> <b>AGE:</b> Gender: Patient ID:	Specimen: Collected: Received: Reported:	

**Immunology**

Test Name	Result	Reference Range	Lab
ANA SCREEN, IFA, W/REFL TITER AND PATTERN			
ANA SCREEN, IFA	NEGATIVE	NEGATIVE	
ANA IFA is a first line screen for detecting the presence of up to approximately 150 autoantibodies in various autoimmune diseases. A negative ANA IFA result suggests ANA-associated autoimmune diseases are not present at this time. Visit Physician FAQs for interpretation of all antibodies in the Cascade, prevalence, and association with diseases at <a href="http://education.QuestDiagnostics.com/faq/FAQ177">http://education.QuestDiagnostics.com/faq/FAQ177</a>			
Physician Comments:			

**PERFORMING SITE:**