

Patient Information	Specimen Information	Client Information		
DOB: AGE: Gender: Phone: Patient ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:			

Test Name INSULIN RESPONSE TO GLUCOSE, 3 SPECIMENS	In Range	Out Of Range	Reference Rang	ge Lab	
TIME 1	FST				
SPECIMEN 1	6.1		uIU/mL		
TIME 2	9:15				
SPECIMEN 2	138.6		uIU/mL		
TIME 3	10:15				
SPECIMEN 3	75.2		uIU/mL		
	Reference Range				
		Fasting:		2.0-19.6	
		30 Minutes	s Post Glucose:	6.0-86.0	
		60 Minutes	s Post Glucose:	8.0-112.0	
		90 Minutes	s Post Glucose:	5.0-68.0	
		120 Minutes	s Post Glucose:	5.0-55.0	
		150 Minutes	s Post Glucose:	3.0-46.0	
		180 Minutes	s Post Glucose:	3.0-20.0	
		240 Minutes	s Post Glucose:	<15.0	
		300 Minutes	s Post Glucose:	<8.0	

Stimulated Insulin reference intervals were established using the Siemens Immulite assay. These values are provided for general guidance only.

This insulin assay shows strong cross-reactivity for some insulin analogs (lispro, aspart, and glargine) and much lower cross-reactivity with others (detemir, glulisine).

## **PERFORMING SITE:**