

| Patient Information | Specimen Information | Client Information |
|---------------------------------------------------------|---------------------------------------------------------------------------------|--------------------|
| DOB: AGE: Gender: Phone: Patient ID: | Specimen: Requisition: Lab Ref #: Collected: Received: Reported: | |

| Test Name | In Range | Out Of Range | Reference Range | Lab |
|------------------------------|-----------|----------------|-----------------|-----|
| HSV 1/2 IGG,TYPE SPECIFIC AB | | | | |
| HSV 1 IGG, TYPE SPECIFIC AB | <0.90 | | index | |
| HSV 2 IGG, TYPE SPECIFIC AB | <0.90 | | index | |
| | Index | Interpretation | | |
| | ----- | ----- | | |
| | <0.90 | Negative | | |
| | 0.90-1.09 | Equivocal | | |
| | >1.09 | Positive | | |

This assay utilizes recombinant type-specific antigens to differentiate HSV-1 from HSV-2 infections. A positive result cannot distinguish between recent and past infection. If recent HSV infection is suspected but the results are negative or equivocal, the assay should be repeated in 4-6 weeks. The performance characteristics of the assay have not been established for pediatric populations, immunocompromised patients, or neonatal screening.

| Patient Information | Specimen Information | Client Information |
|---------------------------------------------------|---------------------------------------------------|--------------------|
| DOB: AGE: Gender: Patient ID: | Specimen: Collected: Received: Reported: | |

Infectious Diseases

| Test Name | Result | Reference Range | Lab |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|-----------------|-----|
| HIV 1/2 ANTIGEN/ANTIBODY, FOURTH GENERATION W/RFL | | | |
| HIV AG/AB, 4TH GEN | NON-REACTIVE | NON-REACTIVE | |
| HIV-1 antigen and HIV-1/HIV-2 antibodies were not detected. There is no laboratory evidence of HIV infection. PLEASE NOTE: This information has been disclosed to you from records whose confidentiality may be protected by state law. If your state requires such protection, then the state law prohibits you from making any further disclosure of the information without the specific written consent of the person to whom it pertains, or as otherwise permitted by law. A general authorization for the release of medical or other information is NOT sufficient for this purpose. For additional information please refer to http://education.questdiagnostics.com/faq/FAQ106 (This link is being provided for informational/educational purposes only.) The performance of this assay has not been clinically validated in patients less than 2 years old. | | | |
| Physician Comments: | | | |

Infectious Diseases

| Test Name | Result | Reference Range | Lab |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|-----------------|-----|
| HEPATITIS C AB W/REFL TO HCV RNA, QN, PCR | | | |
| HEPATITIS C ANTIBODY | NON-REACTIVE | NON-REACTIVE | |
| SIGNAL TO CUT-OFF | 0.02 | <1.00 | |
| HCV antibody was non-reactive. There is no laboratory evidence of HCV infection. In most cases, no further action is required. However, if recent HCV exposure is suspected, a test for HCV RNA (test code 35645) is suggested. For additional information please refer to http://education.questdiagnostics.com/faq/FAQ22v1 (This link is being provided for informational/educational purposes only.) | | | |
| Physician Comments: | | | |

PERFORMING SITE: