

Patient Information	Specimen Information	Client Information
DOB: Gender: Phone: Patient ID: Health ID:	AGE: Fasting: N Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	

COMMENTS: FASTING:NO

Test Name	In Range	Out Of Range	Reference Range	Lab
-----------	----------	--------------	-----------------	-----

CHLAMYDOPHILA PNEUMONIAE

AB (IGG, IGA, IGM)

C. PNEUMONIAE AB (IGG)

C. PNEUMONIAE AB (IGA)

C. PNEUMONIAE AB (IGM)

INTERPRETATION

<1:16

<1:10

PAST INFECTION

1:256 H

titer

titer

titer

REFERENCE RANGE:

IgG <1:64

IgA <1:16

IgM <1:10

The immunofluorescent detection of specific antibodies to Chlamydomphila pneumoniae may be complicated by crossreactive antibodies, non-specific antibody stimulation, or past exposure to similar organisms such as C. psittaci and Chlamydia trachomatis. IgM titers of 1:10 or greater usually indicate recent infection, and any IgG titer may indicate past exposure. IgA is typically present at low titers during primary infection, but may be elevated in recurrent exposures or in chronic infection.

This test was developed and its analytical performance characteristics have been determined by Quest Diagnostics Infectious Disease. It has not been cleared or approved by FDA. This assay has been validated pursuant to the CLIA regulations and is used for clinical purposes.

PERFORMING SITE: