

Patient Information	Specimen Information	Client Information
<b>DOB:</b> <b>AGE:</b> Gender: Phone: Patient ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	

Test Name	In Range	Out Of Range	Reference Range	Lab
COXSACKIE B (1-6) ANTIBODIES, SERUM				
COXSACKIE B1 AB		1:8 H	<1:8	
COXSACKIE B2 AB		1:8 H	<1:8	
COXSACKIE B3 AB		1:8 H	<1:8	
COXSACKIE B4 AB	<1:8		<1:8	
COXSACKIE B5 AB		1:16 H	<1:8	
COXSACKIE B6 AB		1:8 H	<1:8	

**INTERPRETIVE CRITERIA:**

<1:8            Antibody Not Detected  
 > or = 1:8     Antibody Detected

Single titers of > or = 1:32 are indicative of recent infection. Titers of 1:8 or 1:16 may be indicative of either past or recent infection, since CF antibody levels persist for only a few months. A four-fold or greater increase in titer between acute and convalescent specimens confirms the diagnosis. There is considerable crossreactivity among enteroviruses; however the highest titer is usually associated with the infecting serotype.

For more information on this test, go to:  
<http://education.questdiagnostics.com/faq/CoxsackieBvir>  
 uses

This test was developed and its analytical performance characteristics have been determined by Quest Diagnostics Nichols Institute, Chantilly, VA. It has not been cleared or approved by the FDA. This assay has been validated pursuant to the CLIA regulations and is used for clinical purposes.

**PERFORMING SITE:**