

| Patient Information | Specimen Information | Client Information |
|---------------------|----------------------|--------------------|
|                     |                      |                    |
|                     |                      |                    |
|                     |                      |                    |
|                     |                      |                    |
|                     |                      |                    |
|                     |                      |                    |
|                     |                      |                    |
|                     |                      |                    |

## **COMMENTS:**

| Test Name   | In Range   | Out Of Range                        | Reference Range | Lab |
|---|--|-------------------------------------|-----------------|-----|
| COCCIDIOIDES SEROLOGY<br>PANEL, SERUM<br>CF TITER   | <1:2   |                                     |                 |     |
| AB TO F ANTIGEN (IGG), ID<br>AB TO TP ANTIGEN(IGM), ID<br>REFERENCE RANGES:   | NEGATIVE<br>NEGATIVE   |                                     |                 |     |
| CF titer: <1:2<br>AB TO F ANTIGEN (I<br>AB TO TP ANTIGEN (  |  |                                     |                 |     |
| The complement fixation (CF)<br>and its analytical performance<br>have been determined by Quest<br>Infectious Disease. It has not<br>approved by FDA. This assay h<br>pursuant to the CLIA regulat:<br>clinical purposes. | test was dev<br>ce characteri<br>Diagnostics<br>ot been clear<br>nas been vali | reloped<br>stics<br>red or<br>dated |                 |     |