

Patient Name:	Account Number:
Patient Phone:	
Date of Birth (Age):	Account Name:
Sex:	
Referring Dr (NPI #):	Collection Date/Time:
Patient ID:	Received Date/Time:
Specimen ID:	Reported Date/Time:

General Comments and Additional Information

Result Name	Flag	Result	Range/Units	Status	Lab
183194 Chlamydia/GC Amplification					
Chlamydia trachomatis, NAA		Negative	Negative	Final	01
Neisseria gonorrhoeae, NAA		Negative	Negative	Final	01
164099 HSV 1 and 2 Ab, IgG					
HSV 1 IgG, Type Spec		Non Reactive	Non Reactive	Final	02
Please note reference interval change					
HSV-1 IgG testing performed using the Roche Elecsys HSV-1 IgG assay.					
HSV 2 IgG, Type Spec	Abnormal	Reactive	Non Reactive	Final	02
Please note reference interval change					
Current guidelines and recommendations do not recommend routine screening for HSV-2 in asymptomatic individuals, including those that are pregnant. The detection of HSV-2 IgG antibodies in a single sample indicates previous exposure to HSV-2 but does not give information as to the site of HSV infection or the timing of exposure. The predictive value of positive and negative results depends on the population's prevalence and the pretest likelihood of HSV-2. HSV-2 IgG testing					

Patient Name:	Account Number:
Patient Phone:	
Date of Birth (Age):	Account Name:
Sex:	
Referring Dr (NPI #):	Collection Date/Time:
Patient ID:	Received Date/Time:
Specimen ID:	Reported Date/Time:

Result Name	Flag	Result	Range/Units	Status	Lab
-------------	------	--------	-------------	--------	-----

performed using the Roche Elecsys HSV-2 IgG assay.					
012005 RPR, Rfx Qn RPR/Confirm TP					
RPR		Non Reactive	Non Reactive	Final	02
083935 HIV Ab/p24 Ag with Reflex					
HIV Ab/p24 Ag Screen		Non Reactive	Non Reactive	Final	02
HIV-1/HIV-2 antibodies and HIV-1 p24 antigen were NOT detected.					
There is no laboratory evidence of HIV infection.					
HIV Negative					
140659 HCV Antibody					
Hep C Virus Ab		Non Reactive	Non Reactive	Final	02
HCV antibody alone does not differentiate between previously					
resolved infection and active infection. Equivocal and Reactive					
HCV antibody results should be followed up with an HCV RNA test					
to support the diagnosis of active HCV infection.					
006510 HBsAg Screen					
HBsAg Screen		Negative	Negative	Final	02
006734 Hep A Ab, IgM					
Hep A Ab, IgM		Negative	Negative	Final	02
A negative anti-HAV IgM result suggests no recent or current					
HAV infection.					

Patient Name:
Patient Phone:
Date of Birth (Age):
Sex:
Referring Dr (NPI #):
Patient ID:
Specimen ID:

Account Number:
Account Name:
Collection Date/Time:
Received Date/Time:
Reported Date/Time:

Performing Lab

01 - Labcorp Phoenix, 5005 S 40th Street Ste 1200, Phoenix, AZ 85040-2969, (800) 788-9743, Collum, Earle S MD

02 - Labcorp Denver, 8490 Upland Drive, Englewood, CO 80112-7115, (303) 792-2600, Collum, Earle S MD

For Inquiries, the physician may contact the performing lab.

END OF REPORT