

Ordered Items: **Celiac Disease Comprehensive**

Date Collected:	Date Received:	Date Reported:	Fasting: <b>No</b>
-----------------	----------------	----------------	--------------------

**Celiac Disease Comprehensive**

Tests	Result	Flag	Units	Reference Interval
Deamidated Gliadin Abs, IgA <sup>01</sup>	6		units	0-19
<p>Negative 0 - 19                      Weak Positive 20 - 30                      Moderate to Strong Positive &gt;30</p>				
Deamidated Gliadin Abs, IgG <sup>01</sup>	13		units	0-19
<p>Negative 0 - 19                      Weak Positive 20 - 30                      Moderate to Strong Positive &gt;30</p>				
t-Transglutaminase (tTG) IgA <sup>01</sup>	<2			0-3
<p>Negative 0 - 3                      Weak Positive 4 - 10                      Positive &gt;10</p> <p>Tissue Transglutaminase (tTG) has been identified as the endomysial antigen. Studies have demonstrated that endomysial IgA antibodies have over 99% specificity for gluten sensitive enteropathy.</p>				
t-Transglutaminase (tTG) IgG <sup>01</sup>	3		U/mL	0-5
<p>Negative 0 - 5                      Weak Positive 6 - 9                      Positive &gt;9</p>				
Endomysial Antibody IgA <sup>01</sup>	Negative			Negative
Immunoglobulin A, Qn, Serum <sup>01</sup>	311		mg/dL	87-352

**Performing Labs**

**Icon Legend**

▲ Out of reference range    ■ Critical or Alert