

Specimen ID:
Acct #:
Phone: (800) 539-6119

Rte: 00

Control ID:

 Walk-In Lab, LLC
 VART verified

Patient Details
Specimen Details
Physician Details
General Comments & Additional Information
Alternate Control Number:
Alternate Patient ID: Fasting:
Ordered Items

PSA Total+% Free

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
PSA Total+% Free					
Prostate Specific Ag, Serum	9.7	High	ng/mL	0.0-4.0	01
Roche ECLIA methodology. According to the American Urological Association, Serum PSA should decrease and remain at undetectable levels after radical prostatectomy. The AUA defines biochemical recurrence as an initial PSA value 0.2 ng/mL or greater followed by a subsequent confirmatory PSA value 0.2 ng/mL or greater. Values obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.					
PSA, Free	2.46		ng/mL	N/A	01
Roche ECLIA methodology.					
% Free PSA	24.8		%		
The table below lists the probability of prostate cancer for men with non-suspicious DRE results and total PSA between 4 and 10 ng/mL, by patient age (Catalona et al, JAMA 1998, 279:1542).					
	% Free PSA	50-64 yr	65-75 yr		
	0.00-10.00%	56%	55%		
	10.01-15.00%	24%	35%		
	15.01-20.00%	17%	23%		
	20.01-25.00%	10%	20%		
	>25.00%	5%	9%		
Please note: Catalona et al did not make specific recommendations regarding the use of percent free PSA for any other population of men.					

 For inquiries, the physician may contact **Branch: 504-828-2666 Lab: 800-877-5227**