

Specimen ID:
Control ID:

Acct #: 17452095 Phone: (800) 539-6119 Rte: 00
Walk-In Lab, LLC
VART verified
1645 Tiffany Lane
Mandeville, LA 70446

Patient Details

DOB:
Age(y/m/d):
Gender: SSN:
Patient ID:

Specimen Details

Date collected:
Date entered:
Date reported:

Physician Details

Ordering: M ROLLINS
Referring:
ID: 1598746240
NPI: 1598746240

General Comments & Additional Information

Alternate Control Number:
Total Volume: Not Provided

Alternate Patient ID: Not Provided
Fasting: No

Ordered Items

Antineutrophil Cytoplasmic Ab; Venipuncture

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
Antineutrophil Cytoplasmic Ab					
Cytoplasmic (C-ANCA)	<1:20		titer	Neg:<1:20	01
Perinuclear (P-ANCA)	<1:20		titer	Neg:<1:20	01
The presence of positive fluorescence exhibiting P-ANCA or C-ANCA patterns alone is not specific for the diagnosis of Wegener's Granulomatosis (WG) or microscopic polyangiitis. Decisions about treatment should not be based solely on ANCA IFA results. The International ANCA Group Consensus recommends follow up testing of positive sera with both PR-3 and MPO-ANCA enzyme immunoassays. As many as 5% serum samples are positive only by EIA. Ref. AM J Clin Pathol 1999;111:507-513.					
Atypical pANCA	<1:20		titer	Neg:<1:20	01
The atypical pANCA pattern has been observed in a significant percentage of patients with ulcerative colitis, primary sclerosing cholangitis and autoimmune hepatitis.					

01 CB LabCorp Dublin
6370 Wilcox Road, Dublin, OH 43016-1269

Dir: Vincent Ricchiuti, PhD

For inquiries, the physician may contact Branch: 504-828-2666 Lab: 800-282-7300